


# REVOFIL AQUASHINE SOFT FILLER Product Specifications



# Advantages of REVOFIL

- 1. REVOFIL has both biphasic and monophasic physical characteristics introduced by combination of cutting edged peptide technology and crossed linked HA which bring upgraded quality in gel viscosity and elasticity with softer operation.
- 2. REVOFIL contains higher concentration of free HAs, linked with peptide, instead of PBS among cross-linked HA molecules which results in increased stability of HA gel.
- 3. Strong inhibition of endogenous Hyaluronidase increases duration of the gel in skin.
- 4. REVOFIL makes up-regulation of endogenous collagen and hyaluronic acid and to increase skin tightness and lifting.

In overall, peptide technology has endowed higher viscosity and elasticity, increased efficacy of sculpturing with longer lasting, increased gel stability and softer injection of HA gel.




Stay Longer, Stay Active and Stay Spectacular

## Active ingredients

#### Reticulated Hyaluronic acid (23 mg/ml):


Cross-linked hyaluronic acid, non-animal origin Biodegradable gel which is slowly absorbed over the time

#### Oligopeptide-72 (CG-Boostrin):



# REVOLUTIONAL PEPTIDES DERMAL FILLER

#### Elasticity and Fluidity of REVOFIL Fine, Plus, Ultra




#### Rheological Properties

#### Viscoelastic properties of REVOFIL

REVOFIL Ultra, Plus, Fine shows high shear storage modulus (G') and stable shear loss modulus (G") in Dynamic mechanical analysis. The result shows high elasticity, duration and stable fluidity of REVOFIL gels.



# ■ Molecular Structure of reticulated HA gel in REVOFIL



# in-vitro data

#### Dermal / Epidermal Cohesive Effect: Increased 81% Collagen 1, 45% Hyaluronic acid







REVOFIL has shown positive regulation of endogenous collagen and hyaluronic acid synthesis in human fibroblast cell.

#### Direct Inhibition of Hyaluronidase Activity: Decreased 90% Hyaluronidase Activity

#### Interaction of Hyaluronidase and CG-Boostrin




#### Inhibition of Hyaluronidase Activity




REVOFIL contains a strong inhibitor of hyaluronidase. CG-Boostrin, a patented peptide, binds to the enzyme strongly and down-regulates the degradation of reticulated hyaluronic acid which implies increased duration of the gel in skin.

#### Protection of Anti-oxidative stress and Photo-induced Cell Damage

#### Anti-oxidative Stress Recovery




#### Photo-induced Cell death Protection



REVOFIL shows strong anti-oxidant effect against oxidative chemical and photo-damage induced UV in human dermal fibroblasts.

## ■ Decreased Enzymatic Degradation

Concentration of Hyaluronidase: 25unit/ml Incubation temperature: 37°C



REVOFIL showed strong resistance to the enzyme which implies longer lasting in skin compared to competitors. The result indicates that the duration of REVOFIL is much longer than other HA Dermal fillers.

# in-vivo data

#### Clinical Test Data Before and After

One hundred seventy-two (172) patients were injected with REVOFIL Fine (30), REVOFIL Plus (86), REVOFIL Ultra (56), aged  $20\sim56$ . All patients satisfied after injection without allergic effects.

The affirmative results had been reached by 99% in the clinical study. The effectiveness of treatment was expressed in decrease and smoothing of wrinkles in facial area and zone around eyes and increase of volume and improvement in shape of lips, nose and hands.

#### Soft Touch Filler, REUOFIL Fine





Sex : Male Age : Early 60's

Product : REVOFIL Fine Injection Area : Hand Rejuvenation





Sex : Female Age : Early 40's

Product : REVOFIL Fine Injection Area : Upper Eyelids

#### Sub Touch Filler, REDOFIL Plus





Sex : Female Age : Early 40's

Product : REVOFIL Plus Injection Area : Nasolabial Folds





Sex : Female Age : Early 30's

Product : REVOFIL Plus Injection Area : Lip Augmentation

#### Deep Touch Filler, REJOFIL Ultra





Sex : Female Age : Early 30's

Product : REVOFIL Ultra Injection Area : Fore Head





Sex : Female Age : Early 50's

Product: REVOFIL Ultra Injection Area: Nasolabial Folds



# Indication and Summary Table

|                                      | •••••   |   |  |
|--------------------------------------|---|---|--|
|                                      |   |   |  |
|                                      | REVOFIL Fine Soft Touch Filler  | REVOFIL Plus<br>Sub Touch Filler  | REVOFIL Ultra Deep Touch Filler  |
| Concentration of HA                  | 23 mg/mL  | 23 mg/mL  | 23 mg/mL   |
| Needle                               | 30G 1/2   | 27G 1/2   | 27G 1/2  |
| Indication                           | Correction of superficial wrinkles, fine lines, caved scar. Filling of all moderately pronounced wrinkles on the face and tissue. | Filling deep wrinkles and lines.<br>Harmonization of volume and<br>lip contour.   | Redefinition of facial contours.<br>Correction of very deep wrinkles<br>in thick skin tissue.<br>Creation of the volume<br>(Cheeks, chin, temporal area) |
| Injection depth                      | Superficial dermis  | Mid, Deep dermis  | Deep, Subcutaneous dermis  |
| Injection area                       | Tear Troughs<br>Fine lines, Temporal region<br>Upper Cheeks<br>Cheeks, Hollow Cheek   | Fore Head Nose Bridge Nasolabial Folds Marionette lines Tear Troughs Fine lines, Temporal region Upper Cheeks Cheeks, Hollow Cheek Lip Lifting, Augmentation Jaw, Chin, Mandibular region | Fore Head<br>Nose Bridge<br>Nasolabial Folds<br>Marionette lines<br>Cheeks, Hollow Cheek<br>Lip Lifting, Augmentation<br>Jaw, Chin, Mandibular region    |
| Blister pack                         | 1 mL prefilled syringe/pack   | 1 mL prefilled syringe/pack   | 1 mL prefilled syringe/pack  |
| Implant Duration<br>Before touch up* | 10 months on average  | 12 months on average  | 12 months on average   |

<sup>\*</sup> Average duration depends on several factors: Patient's skin type, severity of the wrinkles to be corrected, type of injection and the injected volume.





### CAREGEN CO., LTD.

1027-22, Hogye-dong, Dongan-gu, Anyang-si, Gyeonggi-do 431848 South Korea

**tel**: +82-31-420-9200 **fax**: +82-31-452-3869

e-mail: sales@caregen.com

www.caregen.com / www.renokin.com

# # MEDICA DEPOT

Medica Depot does not claim ownership of the content presented in this product insert - it is a photo reproduction of the product leaflet provided by the Medica Depot team for your convenience. The intellectual property rights, including but not limited to copyrights, trademarks, and other applicable rights, associated with the content of this product insert remain the sole property of the original manufacturer. Medica Depot disclaims any responsibility for any inaccuracies, omissions, or discrepancies in this document. Your use of this product insert is subject to our terms and conditions, available on our website. By accessing and using this document, you acknowledge and agree to the mentioned terms and conditions.